

## tryptophan N-monooxygenase

Cat. No. EXWM-0726

Lot. No. (See product label)

### Introduction

**Description** A heme-thiolate protein (P-450). This enzyme catalyses two successive N-hydroxylations of L-tryptophan, the first steps in the biosynthesis of both auxin and the indole alkaloid phytoalexin camalexin. The product of the two hydroxylations, N,N-dihydroxy-L-tryptophan, is extremely labile and dehydrates spontaneously. The dehydrated product is then subject to a decarboxylation that produces an oxime. It is still not known whether the decarboxylation is spontaneous or catalysed by the enzyme.

**Synonyms** tryptophan N-hydroxylase; CYP79B1; CYP79B2; CYP79B3

### Product Information

**Form** Liquid or lyophilized powder

**EC Number** EC 1.14.13.125

**Reaction** L-tryptophan + 2 O<sub>2</sub> + 2 NADPH + 2 H<sup>+</sup> = (E)-indol-3-ylacetaldoxime + 2 NADP<sup>+</sup> + CO<sub>2</sub> + 3 H<sub>2</sub>O (overall reaction); (1a) L-tryptophan + O<sub>2</sub> + NADPH + H<sup>+</sup> = N-hydroxy-L-tryptophan + NADP<sup>+</sup> + H<sub>2</sub>O; (1b) N-hydroxy-L-tryptophan + O<sub>2</sub> + NADPH + H<sup>+</sup> = N,N-dihydroxy-L-tryptophan + NADP<sup>+</sup> + H<sub>2</sub>O; (1c) N,N-dihydroxy-L-tryptophan = (E)-indol-3-ylacetaldoxime + CO<sub>2</sub> + H<sub>2</sub>O

**Notes** This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

### Storage and Shipping Information

**Storage** Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.