

## Native *Streptomyces griseus* Chitosanase

Cat. No. NATE-0125

Lot. No. (See product label)

### Introduction

**Description** Chitosanase catalyzes the endohydrolysis of  $\beta$  (1,4) linkages between N-acetyl-D-glucosamine and D-glucosamine residues in partially deacetylated chitosan. Chitosanase from *Streptomyces griseus* is capable of hydrolyzing both chitosan and carboxymethyl cellulose. It is used for the lysis of cell walls of fungi belonging to the group Mucorales. It is found in several types of microorganisms.

**Applications** Chitosanase from *Streptomyces griseus* has been used in a study to assess the effect of chitin sources on production of chitinase and chitosanase. Chitosanase from *Streptomyces griseus* has also been used in a study to investigate the effective production of chitinase and chitosanase by *Streptomyces griseus* HUT 6037 using colloidal chitin and various degrees of deacetylation of chitosan. The enzyme from Creative Enzymes has been used for the hydrolysis of the crude cell wall preparation of oomycete *Aphanomyces euteiches*, a major parasite of legume plants. It has also been used for the enzymatic hydrolysis of the fully de-N-acetylated chitosan to get chitosan oligomer mixtures during the preparation of biocompatible chitosan-alginate gel.

**Synonyms** Chitosanase; EC 3.2.1.132; 51570-20-8; Chitosan N-acetylglucosaminohydrolase

### Product Information

**Source** *Streptomyces griseus*

**Form** Lyophilized powder containing potassium phosphate buffer salts

**EC Number** EC 3.2.1.132

**CAS No.** 51570-20-8

**Activity** >50 units/mg protein (Bradford)

**Buffer** H<sub>2</sub>O: soluble, clear to slightly hazy, colorless (10U/0.3mL)

**Unit Definition** One unit will release 1  $\mu$ mole of glucosamine from chitosan per min at pH 5.0 at 37°C as measured in the fluorimetric assay of Osswald, et al. (1992).

### Storage and Shipping Information

**Storage** -20°C