

## pikromycin synthase

Cat. No. EXWM-0784

Lot. No. (See product label)

### Introduction

**Description** A heme-thiolate protein (cytochrome P-450). Involved in the biosynthesis of a number of bacterial macrolide antibiotics containing a desosamine glycoside unit. With narbomycin it hydroxylates at either C-12 to give pikromycin or C-14 to give neopikromycin or both positions to give narvopikromycin. With 10-deoxymethymycin it hydroxylates at either C-10 to give methymycin or C-12 to give neomethymycin or both positions to give novamethymycin.

**Synonyms** PikC; CYP107L1

### Product Information

**Form** Liquid or lyophilized powder

**EC Number** EC 1.14.13.185

**Reaction** (1) narbomycin + NADPH + H<sup>+</sup> + O<sub>2</sub> = pikromycin + NADP<sup>+</sup> + H<sub>2</sub>O; (2) narbomycin + NADPH + H<sup>+</sup> + O<sub>2</sub> = neopikromycin + NADP<sup>+</sup> + H<sub>2</sub>O; (3) narbomycin + 2 NADPH + 2 H<sup>+</sup> + 2 O<sub>2</sub> = novapikromycin + 2 NADP<sup>+</sup> + 2 H<sub>2</sub>O; (4) 10-deoxymethymycin + NADPH + H<sup>+</sup> + O<sub>2</sub> = methymycin + NADP<sup>+</sup> + H<sub>2</sub>O; (5) 10-deoxymethymycin + NADPH + H<sup>+</sup> + O<sub>2</sub> = neomethymycin + NADP<sup>+</sup> + H<sub>2</sub>O; (6) 10-deoxymethymycin + 2 NADPH + 2 H<sup>+</sup> + 2 O<sub>2</sub> = novamethymycin + 2 NADP<sup>+</sup> + 2 H<sub>2</sub>O

**Notes** This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

### Storage and Shipping Information

**Storage** Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.