

cholestanetriol 26-monooxygenase

Cat. No. EXWM-0940

Lot. No. (See product label)

Introduction

Description This mitochondrial cytochrome P-450 enzyme requires adrenodoxin. It catalyses the first three sterol side chain oxidations in bile acid biosynthesis via the neutral (classic) pathway. Can also act on cholesterol, cholest-5-ene-3 β ,7 α -diol, 7 α -hydroxycholest-4-en-3-one, and 5 β -cholestane-3 α ,7 α -diol. The enzyme can also hydroxylate cholesterol at positions 24 and 25. The initial source of the electrons is NADPH, which transfers the electrons to the adrenodoxin via EC 1.18.1.6, adrenodoxin-NADP+ reductase.

Synonyms 5 β -cholestane-3 α ,7 α ,12 α -triol 26-hydroxylase; 5 β -cholestane-3 α ,7 α ,12 α -triol hydroxylase; cholestanetriol 26-hydroxylase; sterol 27-hydroxylase; sterol 26-hydroxylase; cholesterol 27-hydroxylase; CYP27A; CYP27A1; cytochrome P450 27A1'

Product Information

Form Liquid or lyophilized powder

EC Number EC 1.14.15.15

CAS No. 52227-77-7

Reaction 5 β -cholestane-3 α ,7 α ,12 α -triol + 6 reduced adrenodoxin + 6 H⁺ + 3 O₂ = (25R)-3 α ,7 α ,12 α -trihydroxy-5 β -cholestan-26-oate + 6 oxidized adrenodoxin + 4 H₂O (overall reaction); (1a) 5 β -cholestane-3 α ,7 α ,12 α -triol + 2 reduced adrenodoxin + 2 H⁺ + O₂ = (25R)-5 β -cholestane-3 α ,7 α ,12 α ,26-tetraol + 2 oxidized adrenodoxin + H₂O; (1b) (25R)-5 β -cholestane-3 α ,7 α ,12 α ,26-tetraol + 2 reduced adrenodoxin + 2 H⁺ + O₂ = (25R)-3 α ,7 α ,12 α -trihydroxy-5 β -cholestan-26-al + 2 oxidized adrenodoxin + 2 H₂O; (1c) (25R)-3 α ,7 α ,12 α -trihydroxy-5 β -cholestan-26-al + 2 reduced adrenodoxin + 2 H⁺ + O₂ = (25R)-3 α ,7 α ,12 α -trihydroxy-5 β -cholestan-26-oate + 2 oxidized adrenodoxin + H₂O

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.