

## Native Microbial Creatinine Deiminase

Cat. No. NATE-0164

Lot. No. (See product label)

### Introduction

**Description** In enzymology, a creatinine deaminase (EC 3.5.4.21) is an enzyme that catalyzes the chemical reaction: creatinine + H<sub>2</sub>O ↔ N-methylhydantoin + NH<sub>3</sub>. Thus, the two substrates of this enzyme are creatinine and H<sub>2</sub>O, whereas its two products are N-methylhydantoin and NH<sub>3</sub>. This enzyme belongs to the family of hydrolases, those acting on carbon-nitrogen bonds other than peptide bonds, specifically in cyclic amidines. The systematic name of this enzyme class is creatinine iminohydrolase.

**Applications** Creatinine deiminase has been used in a study to assess the application of a creatinine-sensitive biosensor for hemodialysis control. Creatinine deiminase has also been used in a study to investigate the bioelectronic tongue for the simultaneous determination of urea, creatinine and alkaline ions in clinical samples.

**Synonyms** EC 3.5.4.21, creatinine hydrolase; creatinine desiminase; creatinine deaminase; 37289-15-9

### Product Information

<b>Source</b>	Microbial
<b>Form</b>	Lyophilized powder containing mannitol as stabilizer
<b>EC Number</b>	EC 3.5.4.21
<b>CAS No.</b>	37289-15-9
<b>Molecular Weight</b>	mol wt ~260 kDa
<b>Activity</b>	> 25 units/mg protein
<b>Isoelectric point</b>	4.4
<b>pH Stability</b>	pH 7.0 – 11.0 (30°C, 20hr)
<b>Optimum pH</b>	8.5 – 9.5
<b>Thermal stability</b>	Below 65°C (pH 7.5, 1hr)
<b>Optimum temperature</b>	65 – 75°C
<b>Michaelis Constant</b>	3.5 x 10 <sup>-3</sup> M (Creatinine)
<b>Inhibitors</b>	Ag <sup>+</sup> , Hg <sup>++</sup> , o-phenanthroline, monoiodoacetate
<b>Unit Definition</b>	One unit will hydrolyze 1.0 μmole of creatinine to N-methylhydantoin and NH <sub>3</sub> per min at pH 7.5 at 37°C in a coupled system with L-glutamic dehydrogenase.

### Storage and Shipping Information

**Storage** –20°C

