

## L-2-aminoadipate reductase

Cat. No. EXWM-1199

Lot. No. (See product label)

## Introduction

Description This enzyme, characterized from the yeast Saccharomyces cerevisiae, catalyses the reduction of L-2-

aminoadipate to (S)-2-amino-6-oxohexanoate during L-lysine biosynthesis. An adenylation domain activates the substrate at the expense of ATP hydrolysis, and forms L-2-aminoadipate adenylate, which is attached to a peptidyl-carrier protein (PCP) domain. Binding of NADPH results in reductive cleavage of the acyl-S-enzyme intermediate, releasing (S)-2-amino-6-oxohexanoate. Different from EC 1.2.1.31, L-aminoadipate-semialdehyde dehydrogenase, which catalyses a similar transformation in the opposite

direction without ATP hydrolysis.

**Synonyms** LYS2;  $\alpha$ -aminoadipate reductase

## **Product Information**

**Form** Liquid or lyophilized powder

**EC Number** EC 1.2.1.95

Reaction (S)-2-amino-6-oxohexanoate + NADP+ + AMP + diphosphate = L-2-aminoadipate + NADPH + H+ + ATP

 $(overall\ reaction); (1a)\ L-2-aminoadipyl-[LYS2\ peptidyl-carrier-protein] + AMP + diphosphate = L-2-aminoadipate + holo-[LYS2\ peptidyl-carrier-protein] + ATP; (1b)\ (S)-2-amino-6-oxohexanoate + holo-[LYS2\ peptidyl-carrier-protein] + ATP; (1b)\ (S)-2-amino-6-oxohexanoate + holo-[LYS2\ peptidyl-carrier-protein] + ATP; (1b)\ (S)-2-amino-6-oxohexanoate + holo-[LYS2\ peptidyl-carrier-protein] + ATP; (1c)\ (S)-2-amino-6-oxohexanoate + holo-[L$ 

 $peptidyl-carrier-protein] + NADP+ = L-2-aminoadipyl-[LYS2\ peptidyl-carrier-protein] + NADPH + H+$ 

**Notes** This item requires custom production and lead time is between 5-9 weeks. We can custom produce

according to your specifications.

## Storage and Shipping Information

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.

**Tel:** 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com

1/1