

Native Human Elastase

Cat. No. NATE-0213

Lot. No. (See product label)

Introduction

Description Neutrophil elastase is a serine proteinase in the same family as chymotrypsin and has broad substrate

specificity. Secreted by neutrophils and macrophages during inflammation, it destroys bacteria and host tissue. It also localizes to Neutrophil extracellular traps (NETs), via its high affinity for DNA, an unusual

property for serine proteases.

Applications Elastase from Creative Enzymes has been used to digest fibronectin. The results were compared with

fibronectin digestion by crude human leuk ocyte homogenateto examine the presence of fibronectin peptides in saliva of patients with Sjögren's syndrome. It has also been used as a reference to determine the elastase activity in cell lysates. This study examined the effect of all-trans retinoic acid on pr ocoagulant and fibrinolytic activities of cultured blast cells. These blast cells were from patients with acute promyel ocytic leukemia. Elastase from human leuk ocytes has been used in a study that

determined that fragments of Nle3-angiotensin (1-7) accelerate healing in dermal models.

Synonyms ELANE; elastase; EC 3.4.21.37; leukocyte elastase; ELA2; elastase 2; neutrophil elaszym; serine elastase;

lysosomal elastase; neutrophil elastase; polymorphonuclear leukocyte elastase; elaszym; granulocyte

elastase

Product Information

Species Human

Source Human leuk ocytes

Form Lyophilized from 0.05 M sodium acetate (pH 5.5) and 0.6 M NaCl

EC Number EC 3.4.21.37

CAS No. 9004-06-2

Molecular

29 kDa

Weight

Activity > 50 units/mg protein (Bradford)

Isoelectric

point

8.77-9.55

Pathway Activation of Matrix Metalloproteinases, organism-specific biosystem; C-MYB transcription factor network,

organism-specific biosystem; Degradation of the extracellular matrix, organism-specific biosystem; Extracellular matrix organization, organism-specific biosystem; Systemic lupus erythematosus, organism-specific biosystem; Systemic lupus erythematosus, conserved biosystem; Transcriptional misregulation

in cancer, organism-specific biosystem

Function bacterial cell surface binding; cytokine binding; endopeptidase activity; heparin binding; peptidase

activity; protease binding; protein binding; serine-type endopeptidase activity

Unit One unit will release one nanomole of p-nitrophenol per sec from Bec-L-alanine p-nitrophenyl ester at pH

Definition 6.5 at 37°C.

Storage and Shipping Information

Storage –20°C

Tel: 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com 1/2