

4,4'-diapophytoene desaturase (4,4'-diapolycopene-forming)

Cat. No. EXWM-1409

Lot. No. (See product label)

Introduction

Description The enzyme catalyses four successive dehydrogenations, resulting in production of 4,4'-diapolycopene. While the enzyme from *Staphylococcus aureus* was only shown to produce 4,4'-diaponeurosporene in vivo, it is able to catalyse the last reaction in vitro.

Synonyms dehydrosqualene desaturase (ambiguous); CrtN (ambiguous); 4,4'-diapophytoene:FAD oxidoreductase (ambiguous); 15-cis-4,4'-diapophytoene:FAD oxidoreductase; 4,4'-diapophytoene desaturase (ambiguous)

Product Information

Form Liquid or lyophilized powder

EC Number EC 1.3.8.2

Reaction 15-cis-4,4'-diapophytoene + 4 FAD = all-trans-4,4'-diapolycopene + 4 FADH₂ (overall reaction); (1a) 15-cis-4,4'-diapophytoene + FAD = all-trans-4,4'-diapophytofluene + FADH₂; (1b) all-trans-4,4'-diapophytofluene + FAD = all-trans-4,4'-diapo- ζ -carotene + FADH₂; (1c) all-trans-4,4'-diapo- ζ -carotene + FAD = all-trans-4,4'-diaponeurosporene + FADH₂; (1d) all-trans-4,4'-diaponeurosporene + FAD = all-trans-4,4'-diapolycopene + FADH₂

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.