

Hck from Human, Recombinant

Cat. No. NATE-0335

Lot. No. (See product label)

Introduction

Description Hck is a member of the Src family of non-receptor associated protein tyrosine kinases (PTK). Hck has been shown to associate with and be activated by Bcr-Abl, which is a protein-tyrosine kinase that is expressed in chronic myelogenous leukemia.

Applications Hck is a protein-tyrosine kinase that is expressed in hemopoietic cells. It is used in leukemia research

Synonyms Hck; HCK; tyrosine-protein kinase HCK; JTK9; p59Hck; p61Hck

Product Information

Species Human

Source E. coli BL21

Form Supplied as a solution in 50 mM Tris HCl, pH 7.4, 1 mM EDTA and 1 mM DTT

Activity > 200 units

Pathway Alpha-synuclein signaling, organism-specific biosystem; B Cell Receptor Signaling Pathway, organism-specific biosystem; CXCR4-mediated signaling events, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class I PI3K signaling events, organism-specific biosystem; Cytokine Signaling in Immune system, organism-specific biosystem

Function ATP binding; non-membrane spanning protein tyrosine kinase activity; nucleotide binding; protein binding; protein tyrosine kinase activity

Unit Definition One unit will phosphorylate 1.0 picomole of Tyr15 of the peptide substrate pcdc2 6-20 per minute at 30°C at pH 7.4.

Usage and Packaging

Package vial of > 200 units

Storage and Shipping Information

Storage -70°C