

Native Human Lysozyme

Cat. No. NATE-0433

Lot. No. (See product label)

Introduction

Description Lysozymes, also known as muramidase or N-acetylmuramide glycanhydrolase, are glycoside hydrolases. These are enzymes (EC 3.2.1.17) that damage bacterial cell walls by catalyzing hydrolysis of 1,4-beta-linkages between N-acetylmuramic acid and N-acetyl-D-glucosamine residues in a peptidoglycan and between N-acetyl-D-glucosamine residues in chitodextrins. Lysozyme is abundant in a number of secretions, such as tears, saliva, human milk, and mucus. It is also present in cytoplasmic granules of the macrophages and the polymorphonuclear neutrophils (PMNs). Large amounts of lysozyme can be found in egg white. C-type lysozymes are closely related to alpha-lactalbumin in sequence and structure, making them part of the same family. In humans, the lysozyme enzyme is encoded by the LYZ gene.

Synonyms muramidase; globulin G; mucopolysaccharide glucohydrolase; globulin G1; N,O-diacetylmuramidase; lysozyme g; L-7001; 1,4-N-acetylmuramidase; mucopolysaccharide N-acetylmuramoylhydrolase; PR1-lysozyme; lysozyme; LYZ; LYM; EC 3.2.1.17; 9001-63-2

Product Information

Species	Human
Source	Human neutrophils
Form	Lyophilized from 50 mM sodium acetate, pH 6.0, with 100 mM NaCl
EC Number	EC 3.2.1.17
CAS No.	9001-63-2
Purity	> 95% (SDS-PAGE)
Activity	30,000 Sugar units per mg
Pathway	Amyloids, organism-specific biosystem; C-MYB transcription factor network, organism-specific biosystem; Disease, organism-specific biosystem; Salivary secretion, organism-specific biosystem; Salivary secretion, conserved biosystem
Function	hydrolase activity, acting on glycosyl bonds; lysozyme activity
Unit Definition	One unit is defined as the amount of enzyme that digest powdered cells of <i>Micrococcus lysodeikticus</i> , causing a decrease in absorbance of 0.001 per minute at 37°C, pH 7.0.

Storage and Shipping Information

Storage -20°C