

asparaginyl-tRNA synthase (glutamine-hydrolysing)

Cat. No. EXWM-5809

Lot. No. (See product label)

Introduction

Description This reaction forms part of a two-reaction system for producing asparaginyl-tRNA in *Deinococcus radiodurans* and other organisms lacking a specific enzyme for asparagine synthesis. In the first step, a non-discriminating ligase (EC 6.1.1.23, aspartate-tRNAAsn ligase) mischarges tRNAAsn with aspartate, leading to the formation of Asp-tRNAAsn. The aspartyl-tRNAAsn is not used in protein synthesis until the present enzyme converts it into asparaginyl-tRNAAsn (aspartyl-tRNAAsp is not a substrate for this reaction). Ammonia or asparagine can substitute for the preferred substrate glutamine.

Synonyms Asp-AdT; Asp-tRNAAsn amidotransferase; aspartyl-tRNAAsn amidotransferase; Asn-tRNAAsn:L-glutamine amido-ligase (ADP-forming); aspartyl-tRNAAsn:L-glutamine amido-ligase (ADP-forming)

Product Information

Form Liquid or lyophilized powder

EC Number EC 6.3.5.6

CAS No. 37211-76-0

Reaction $\text{ATP} + \text{L-aspartyl-tRNAAsn} + \text{L-glutamine} + \text{H}_2\text{O} = \text{ADP} + \text{phosphate} + \text{L-asparaginyl-tRNAAsn} + \text{L-glutamate}$

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.