

Transglutaminase from Mouse, Recombinant

Cat. No. NATE-1727

Lot. No. (See product label)

Introduction

Description This enzyme belongs to the family of transferases, specifically those transferring phosphorus-containing groups (phosphotransferases) with a phosphate group as acceptor.

Applications Recombinant mouse transglutaminase is a homodimer (α_2) composed of two chains held together by non covalent bonds. After activation of the zymogen by Thrombin and Ca^{2+} to its active form (α_2^* , Factor XIIIa), Factor XIIIa catalyzes the formation of covalent bridges (ϵ - γ -glutamyl) lysine bonds between fibrin units to increase the elasticity of the clot network. The resulting cross-linked fibrin is insoluble and resistant to lysis.

Synonyms transglutaminase; EC 2.3.2.13; 80146-85-6; transglutaminase; Factor XIIIa; fibrinolygase; fibrin stabilizing factor; glutaminylopeptide γ -glutamyltransferase; polyamine transglutaminase; tissue transglutaminase; R-glutaminylopeptide:amine γ -glutamyl transferase; protein-glutamine γ -glutamyltransferase; TG1

Product Information

Species Mouse

Source Insect cells

Appearance Liquid

Form The transglutaminase is formulated in 10 mM Sodium Phosphate pH 8.0, 15 mM NaCl. Sample contains 50% glycerol. Biotinylated-transglutaminase is a Ca^{2+} -dependent enzyme.

EC Number EC 2.3.2.13

CAS No. 80146-85-6

Molecular Weight 83 kDa

Purity > 95% (visually by SDS-PAGE)

Activity > 750 U/mg [Activity is determined by measuring the rate of fluorescence enhancement after transglutaminase-catalyzed monodansylcadaverine-incorporation into N,N-dimethylated casein according to Lorand et al., Anal. Biochem. 44 (221-231)].

Activators Add Thrombin and 10 mM Ca^{2+} to activate Transglutaminase.

Unit Definition 1 U is defined as the increase in fluorescence intensity of 1 a.u./min (measured on a Cary eclipse fluorescence spectrophotometer, Varian; $\lambda_{\text{ex}} = 332 \text{ nm}$, $\lambda_{\text{em}} = 500 \text{ nm}$; band filter = 5 nm; detector strength = 600 V; temperature = 37 °C, assay volume = 1 ml).

Usage and Packaging

Package 50 μg

Storage and Shipping Information

Storage Store working aliquots at $\leq -20^\circ\text{C}$. Avoid repeated freezing and thawing.