

Formate Dehydrogenase (Crude Enzyme)

Cat. No. NATE-1799

Lot. No. (See product label)

Introduction

Description Formate Dehydrogenases are a set of enzymes that catalyse the oxidation of formate to carbon dioxide, donating the electrons to a second substrate, such as NAD + in formate:NAD + oxidoreductase (EC 1. 2. 1. 2) or to a cytochrome in formate:ferricytochrome-b1 oxidoreductase (EC 1. 2. 2. 1). This product with the indicated enzyme activity was briefly purified from engineered E. coli.

Applications synthesis; biotechnology; medicine; analysis

Synonyms formate-NAD oxidoreductase; FDH I; FDH II; N-FDH; formic hydrogen-lyase; formate hydrogenlyase; hydrogenlyase; NAD-linked formate dehydrogenase; NAD-dependent formate dehydrogenase; formate dehydrogenase (NAD); NAD-formate dehydrogenase; formate benzyl-viologen oxidoreductase; formic acid dehydrogenase

Product Information

Source E. coli

Appearance Clear to translucent yellow solution

EC Number EC 1.2.1.2

CAS No. 9028-85-7

Activity Undetermined

Reaction formate + NAD + = CO₂ + NADH

Notes Since this product needs to be freshly prepared, it will take about 2 weeks after you confirm the order. Each time of the freeze-thawing may cause partial inactivation. Therefore, it should be dispensed as required and stored at -20 ° C or lower. With the preservation of the extension of time, the enzyme activity will decline to a certain extent, so the product should be used as soon as possible. This product may have turbidity or precipitation in the production and preservation process, it can be mixed after melting and will not affect the normal use. This product is limited to scientific research use, shall not be used for clinical diagnosis or treatment, shall not be used for food or medicine, shall not be stored in ordinary residential. For your safety and health, please wear an experimental suit and wear disposable gloves.

Usage and Packaging

Package 100ml

Storage and Shipping Information

Storage at -20 °C or lower, for at least 1 month.