

## Native Rat Protein Kinase C

Cat. No. NATE-0573

Lot. No. (See product label)

### Introduction

**Description** Protein kinase C is a family of protein kinase enzymes that are involved in controlling the function of other proteins through the phosphorylation of hydroxyl groups of serine and threonine amino acid residues on these proteins. PKC enzymes in turn are activated by signals such as increases in the concentration of diacylglycerol (DAG) or calcium ions (Ca<sup>2+</sup>). Hence PKC enzymes play important roles in several signal transduction cascades.

**Synonyms** EC 2.7.11.13; 141436-78-4; Protein kinase C; PKC; calcium-dependent protein kinase C; calcium-independent protein kinase C; calcium/phospholipid dependent protein kinase; cPKC $\alpha$ ; cPKC $\beta$ ; cPKC $\gamma$ ; nPKC $\delta$ ; nPKC $\epsilon$ ; nPKC $\eta$ ; nPKC $\theta$ ; PKC $\alpha$ ; PKC $\beta$ ; PKC $\gamma$ ; PKC $\delta$ ; PKC $\epsilon$ ; PKC $\zeta$ ; Pkc1p; protein kinase C $\epsilon$ ; STK24

### Product Information

**Species** Rat

**Source** Rat brain

**Form** Type I, buffered aqueous glycerol solution; Type II, lyophilized powder.

**EC Number** EC 2.7.1.37

**CAS No.** 141436-78-4

**Molecular Weight** mol wt 82 kDa by SDS-PAGE

**Purity** > 90% (SDS-PAGE)

**Activity** Type I, 50-200 units/mL

**Buffer** Solution in 50% glycerol containing 20 mM Tris, pH 7.5, 0.5 mM EDTA, 0.5 mM EGTA, 5 mM DTT, 100 mM NaCl, 0.02% Tween 20, and 1  $\mu$ g/ml leupeptin.

**Unit Definition** One unit will transfer 1 nmol of phosphate per min from ATP to histone H1 at pH 7.4 at 30°C.

### Storage and Shipping Information

**Stability** -70°C