

## Native Elizabethkingia miricola PNGase F

Cat. No. NATE-0602

Lot. No. (See product label)

### Introduction

**Description** In enzymology, a peptide-N4-(N-acetyl-beta-glucosaminy) asparagine amidase (EC 3.5.1.52) is an enzyme that catalyzes a chemical reaction that cleaves a N4-(acetyl-beta-D-glucosaminy)asparagine residue in which the glucosamine residue may be further glycosylated, to yield a (substituted) N-acetyl-beta-D-glucosaminyamine and a peptide containing an aspartate residue. This enzyme belongs to the family of hydrolases, specifically those acting on carbon-nitrogen bonds other than peptide bonds in linear amides.

**Applications** Used to deglycosylate protein.

**Synonyms** glycopeptide N-glycosidase; glycopeptidase; N-oligosaccharide glycopeptidase; N-glycanase; glycopeptidase; Jack-bean glycopeptidase; PNGase A; PNGase F; glycopeptide N-glycosidase; peptide-N4-(N-acetyl-β-glucosaminy)asparagine amidase; EC 3.5.1.52; PNGase F; 83534-39-8

### Product Information

**Source** Elizabethkingia miricola

**Form** Supplied as a solution in 20 mM Tris HCl, pH 7.5, 50 mM NaCl and 1 mM EDTA

**EC Number** EC 3.5.1.52

**CAS No.** 83534-39-8

**Molecular Weight** 36 kDa

**Activity** > 20,000 units/mg protein and > 5,000 units/ml

**pH Stability** active in the pH range of 6-10

**Optimum pH** 8.6

**Unit Definition** One unit will catalyze the release of N-linked oligosaccharides from 1 nanomole of denatured ribonuclease B in one minute at 37°C at pH 7.5 monitored by SDS-PAGE. One Sigma unit of PNGase F activity is equal to 1 IUB milliunit.

### Storage and Shipping Information

**Storage** 2-8°C

**Stability** The product remains active for at least 12 months when stored properly. Exposure for several days to ambient temperatures will not reduce activity. Do Not Freeze.