

Chemically modified *Pseudomonas* species Lipoprotein Lipase

Cat. No. DIA-282

Lot. No. (See product label)

Introduction

Description Enzyme that hydrolyzes triglycerides into three free fatty acids and glycerol. Take advantage of the enhanced liquid stability of this enzyme. Rely on the proven diagnostic quality of this product.

Applications Use Lipoprotein lipase in diagnostic tests for the determination of triglycerides together with Glycerol Kinase and Glycerol-3-phosphate Dehydrogenase.

Synonyms Lipoprotein lipase; LPL; Clearing factor lipase; Diacylglycerol lipase; Diglyceride lipase

Product Information

Source *Pseudomonas* species

Appearance Brownish lyophilizate

Molecular Weight 47 kD

Activity >10 U/mg lyophilizate

Contaminants ATPase: <0.005 Catalase: <1.0 Glycerokinase: <0.001 Hexokinase: <0.005 "NADH oxidase": <0.001 Uricase: <0.005

pH Stability 6.0-10.0

Optimum pH 7.5

Thermal stability Up to +50°C

Specificity Lipoprotein Lipase has both lipolytic and sterol ester hydrolytic activities. It hydrolyzes triacylglycerols in chylomicrons, lipoproteins and diacylglycerols. With human plasma as substrate triglycerides are hydrolyzed more rapidly than cholesterol esters. The effects of pH and ionic strength on the enzymatic activity are somewhat different between the hydrolysis of triglyceride and of cholesterol ester depending on the different states of these substrates in the plasma or the transfer of the reaction products at the interface of substrates.

Storage and Shipping Information

Stability At +2 to +8°C within specification range for 12 months. Store dry.