

## β-Xylosidase from Lactobacillus brevis, Recombinant

Cat. No. NATE-1191

Lot. No. (See product label)

### Introduction

**Description** Releases reducing sugars from birchwood xylan (X0502), also catalyzes the hydrolysis of 4-methylumbelliferyl-β-D-cellobioside and 4-methylumbelliferyl-β-D-glucopyranoside. This enzyme does not possess endo-xylanase, arabinoxylanase or β-glucanase activities. β-Xylosidase undergoes post-translation glycosylation which has been shown to be critical for its proper activity and stability. Deglycosylation altered the the optimum temperature and pH for activity and decreased its thermostability.

**Synonyms** β-Xylosidase; β-Xylosidase, thermostable; 9025-53-0

### Product Information

**Source** Lactobacillus brevis ATCC 367

**Form** Supplied in 3.2 M ammonium sulphate

**EC Number** EC 3.2.1.37

**CAS No.** 9025-53-0

**Molecular Weight** 63608.3 Da

**Purity** > 95 % as judged by SDS-PAGE

**Activity** 26.76 U/mg

**Concentration** 210.83 U/ml

**Optimum pH** 6

**Optimum temperature** 40°C

**Unit Definition** One unit is defined as the amount of enzyme required to release 1μmol of oNP per minute from oNP-β-D-xylopyranoside (5 mM) in 20 mM sodium phosphate buffer, pH 6.0, at 35°C, and using an extinction coefficient of 18000 M<sup>-1</sup>cm<sup>-1</sup>.

### Storage and Shipping Information

**Storage** Store at 4°C (shipped at room temperature)