

Xylosidase 43A from Bacteroides ovatus, Recombinant

Cat. No. NATE-1527

Lot. No. (See product label)

Introduction

Description Releases reducing sugars from birchwood xylan (X0502), also catalyzes the hydrolysis of 4-methylumbelliferyl- β -D-cellobioside and 4-methylumbelliferyl- β -D-glucopyranoside. This enzyme does not possess endo-xylanase, arabinoxylanase or β -glucanase activities. β -Xylosidase undergoes post-translation glycosylation which has been shown to be critical for its proper activity and stability. Deglycosylation altered the the optimum temperature and pH for activity and decreased its thermostability.

Synonyms β -Xylosidase; EC 3.2.1.37; 9025-53-0; Xylosidase

Product Information

Species	Bacteroides ovatus
Source	E. coli
Form	35 mM NaHepes buffer, pH 7.5, 750 mM NaCl, 200 mM imidazol, 3.5 mM CaCl ₂ , 0.02% sodium azide and 25% (v/v) glycerol
EC Number	EC 3.2.1.37
CAS No.	9025-53-0
Molecular Weight	39.3 kDa
Purity	>90% as judged by SDS-PAGE
Concentration	1 mg/mL
Optimum pH	6.8
Optimum temperature	37 °C
Specificity	Xylooligosaccharides

Storage and Shipping Information

Storage This enzyme is shipped at room temperature but should be stored at -20 °C.