

Pectate Lyase from *Pectobacterium carotovorum*, Recombinant

Cat. No. NATE-1559

Lot. No. (See product label)

Introduction

Description Pectate lyase (EC 4.2.2.2) is an enzyme involved in the maceration and soft rotting of plant tissue. Pectate lyase is responsible for the eliminative cleavage of pectate, yielding oligosaccharides with 4-deoxy- α -D-mann-4-enuronosyl groups at their non-reducing ends. The protein is maximally expressed late in pollen development. It has been suggested that the pollen expression of pectate lyase genes might relate to a requirement for pectin degradation during pollen tube growth. This enzyme belongs to the family of lyases, specifically those carbon-oxygen lyases acting on polysaccharides.

Synonyms (1->4)-alpha-D-galacturonan lyase; polygalacturonic transeliminase; pectic acid transeliminase; polygalacturonate lyase; endopectin methyltranseliminase; pectate transeliminase; endogalacturonate transeliminase; pectic acid lyase; pectic lyase; alpha-1,4-D-endopolygalacturonic acid lyase; PGA lyase; PPase-N; endo-alpha-1,4-polygalacturonic acid lyase; polygalacturonic acid lyase; pectin transeliminase; Polygalacturonic acid trans-eliminase; Pectate lyase; EC 4.2.2.2

Product Information

Species	Pectobacterium carotovorum
Source	E. coli
Form	35 mM NaHepes buffer, pH 7.5, 750 mM NaCl, 200 mM imidazol, 3.5 mM CaCl ₂ , 0.02% sodium azide and 25% (v/v) glycerol
EC Number	EC 4.2.2.2
CAS No.	9015-75-2
Molecular Weight	63.6 kDa
Purity	>90% as judged by SDS-PAGE
Concentration	1 mg/mL
Optimum pH	8.5
Optimum temperature	37 °C
Specificity	Polygalacturonans

Storage and Shipping Information

Storage This enzyme is shipped at room temperature but should be stored at -20 °C.