

## Alcohol dehydrogenase from Equine, Recombinant

Cat. No. NATE-1584

Lot. No. (See product label)

### Introduction

**Description** Alcohol dehydrogenases (ADH) are a group of dehydrogenase enzymes that occur in many organisms and facilitate the interconversion between alcohols and aldehydes or ketones with the reduction of nicotinamide adenine dinucleotide (NAD<sup>+</sup> to NADH). In Humans and many other animals, they serve to break down alcohols that otherwise are toxic, and they also participate in generation of useful aldehyde, ketone, or alcohol groups during biosynthesis of various metabolites. In yeast, plants, and many bacteria, some alcohol dehydrogenases catalyze the opposite reaction as part of fermentation to ensure a constant supply of NAD<sup>+</sup>.

**Synonyms** aldehyde reductase; ADH; alcohol dehydrogenase (NAD); aliphatic alcohol dehydrogenase; ethanol dehydrogenase; NAD-dependent alcohol dehydrogenase; NAD-specific aromatic alcohol dehydrogenase; NADH-alcohol dehydrogenase; NADH-aldehyde dehydrogenase; primary alcohol dehydrogenase; yeast alcohol dehydrogenase; EC 1.1.1.1

### Product Information

**Species** Equine

**Source** E. coli

**EC Number** EC 1.1.1.1

**CAS No.** 9031-72-5

**Activity** >0.5 U/mg

**Unit Definition** 1 U corresponds to the amount of enzyme which reduces 1  $\mu$ mol benzaldehyde per minute at pH 7.0 and 30 °C.

### Storage and Shipping Information

**Storage** Store at -20°C