

Human Factor XIII

Cat. No. CZY-010

Lot. No. (See product label)

Introduction

Description Factor XIII is the zymogenic form of the glutamyl-peptide g-glutamyl transferase factor XIIIa (fibrinolygase, plasma transglutaminase, fibrin stabilizing factor, E.C. 2.3.2.13). Factor XIII is unique among transamidases in that it is a zymogen in vivo. Factor XIII is found both extracellularly in plasma and intracellularly in platelets, megakaryocytes, monocytes, placenta, uterus, liver and prostrate tissues. Plasma factor XIII is synthesized in the liver and circulates as a tetramer ($M_r=320,000$), composed of 2 pairs of nonidentical subunits (A₂B₂). The intra-cellular forms are synthesized in the tissues where they reside as dimers ($M_r=146,000$) of 2 identical A chains (A₂). The A subunits of plasma and intracellular forms of factor XIII are functionally identical. The A subunit contains 6 free sulfhydryl groups one of which is the active site. The concentration of factor XIII in plasma (A₂B₂) is approximately 30 mg/ml. It is the last of the zymogens to become activated in the coagulation cascade and it is the only enzyme in this system that is not a serine protease. The conversion of plasma factor XIII (A₂B₂) to the active transamidase factor XIIIa (A₂') results from hydrolysis of the Arg36-Gly37 at the NH₂-terminus of the A subunit by thrombin. Full expression of activity is achieved only after the Ca²⁺ ($K_d=10^{-3}M$) and fibrin(ogen) ($K_d=10^{-8}M$) dependent dissociation of the B subunit dimer from the A₂' dimer. In the coagulation cascade, factor XIIIa functions to stabilize the fibrin clot by crosslinking the a and g-chains of fibrin. Other proteins known to be substrates for Factor XIIIa which may be hemostatically important include fibronectin, α_2 -antiplasmin, collagen, factor V, von Willebrand Factor and thrombospondin. Factor XIII is purified from fresh frozen human plasma by a modification of the procedures described by Folke and Lorand involving barium citrate, ammonium sulfate and glycine precipitations, ion exchange chromatography and gel filtration. Factor XIII is homogeneous as judged by SDS PAGE, with a specific activity of ~50 units/mg. Factor XIII is supplied in 50% glycerol containing 0.5 mM EDTA, for storage at -20°C.

Product Information

Source	Human
Formulation	50% (vol/vol) glycerol/500uM EDTA
CAS No.	9013-56-3
Molecular Weight	320000
Purity	>95% by SDS-PAGE
Specific Activity	91 U/mg
Concentration	9.1 mg/mL
Isoelectric point	5.2
Structure	tetramer (A ₂ B ₂) of noncovalently associated pairs of 2 non identical subunits, A ($M_r=75,000$), B ($M_r=88,000$); the A subunit has 6 free sulfhydryls and the potential active site
Buffer	50% Glycerol / 500 uM EDTA (vol/vol)
Localization	Plasma
Extinction	13.8

Extraction coefficient 15.0

Percent carbohydrate A chain: 1%; B chain: 5%

Usage and Packaging

Package 100 µg

Storage and Shipping Information

Storage -20°C

Stability 12 months