

## Human Protein C

Cat. No. CZY-017

Lot. No. (See product label)

### Introduction

**Description** The vitamin K-dependent zymogen, protein C, is synthesized in the liver as a single chain polypeptide and is subsequently converted to a disulfide linked heterodimer, by removal of a dipeptide (Lys-146 and Arg-147) from the precursor molecule. Trace quantities of the single chain form have been observed in plasma. The light chain, which is responsible for the calcium dependent binding of protein C to phospholipid vesicles, contains 11  $\gamma$ -carboxyglutamic acid (gla) residues, 1  $\beta$ -hydroxyaspartic acid residue, and 2 epidermal growth factor (EGF) homology domains. The serine protease catalytic triad is located in the heavy chain. Human protein C is susceptible to proteolytic cleavage of a peptide (Mr=3000) from the COOH-terminal end of the heavy chain, yielding an altered form referred to as  $\beta$ -protein C. No functional distinction between  $\alpha$ - and  $\beta$ -protein C has been observed. A single cleavage at Arg-12 (Arg-14 in bovine) of the heavy chain of human protein C converts the zymogen into the serine protease, activated protein C. This cleavage is catalyzed by a complex between  $\alpha$ -thrombin and the endothelial cell surface protein thrombomodulin. In contrast to the other vitamin K dependent coagulation factors, activated protein C functions as an anticoagulant by catalyzing the proteolytic inactivation of factors Va and VIIIa. APC also contributes to the fibrinolytic response by complex formation with plasminogen activator inhibitors. Bovine protein C is prepared from fresh citrated bovine plasma by a modification of the Walker procedure, as described by Haley et al. Human protein C is prepared from fresh frozen citrated human plasma using a combination of immunoaffinity chromatography, and conventional techniques. Protein C is provided in 50% (vol/vol) glycerol/H<sub>2</sub>O and should be stored at -20°C. Purity is determined by SDS-PAGE analysis and activity is measured using a chromogenic substrate based assay.

### Product Information

<b>Source</b>	Human
<b>Formulation</b>	50% glycerol/water (v/v)
<b>CAS No.</b>	42617-41-4
<b>Molecular Weight</b>	62000
<b>Purity</b>	>95% by SDS-PAGE
<b>Specific Activity</b>	<1% APC Activity
<b>Concentration</b>	7.6 mg/mL
<b>Isoelectric point</b>	4.4-4.8
<b>Structure</b>	two chains, Mr=41,000 and 21,000, disulfide linked, NH <sub>2</sub> -terminal gla domain two EGF domains
<b>Buffer</b>	50% Glycerol/H <sub>2</sub> O (v/v)
<b>Localization</b>	Plasma
<b>Extinction coefficient</b>	14.5
<b>Percent</b>	0.23

**Percent carbohydrate** 0.25

**Post-translational modifications** nine gla residues one  $\beta$ -hydroxyaspartate

### **Usage and Packaging**

**Package** 100  $\mu$ g

### **Storage and Shipping Information**

**Storage** -20°C

**Stability** 12 months